

Dynamics Notes

1 – Newton's Laws

Newton's 1st Law:

An object *in motion will stay in motion* and an object *at rest will stay at rest* unless... *an outside force is applied*

aka. Law of Inertia

Newton's 2nd Law:

The acceleration of a body is proportional to the force and inversely proportional to the mass...
As a formula:

$$F_{\text{net}} = ma$$

Newton's 3rd Law:

For every *action force* there is an *equal and opposite reaction force*


As a formula:

$$F_{\text{action}} = -F_{\text{reaction}}$$


Free Body Diagrams: (Draw one for **EVERY** force question)

- 1) Represent the object as a box and,
- 2) Represent all forces as arrows
 - The arrows are vectors so their direction is critical
 - Start in the center and draw outwards
 - The size of the arrow represents the magnitude of the force


1. A puck slides along frictionless ice.



2. A dragster accelerates from rest.



3. A block of wood slides down an incline



Ex. A student pulls straight upwards with a force of 650 N on their 15 kg backpack. What is the backpack's acceleration?



$$F_{\text{net}} = F_{\text{app}} - F_g = ma$$

$$a = \frac{F_{\text{app}} - F_g}{m}$$

$$= \frac{650\text{N} - 147\text{N}}{15\text{kg}}$$


$$F_g = mg$$

$$= (15\text{kg})(9.8\text{m/s}^2)$$

$$= 147\text{N}$$

$$= \boxed{34\text{ m/s}^2}$$

Ex. A 1200kg car accelerates at 5.85 m/s². If the coefficient of friction acting on the car is roughly 0.24, how much force does the engine exert?



$$F_f = \mu F_N$$

only true on level surface

$$= \mu F_g$$

$$= \mu mg$$

$$= (0.24)(1200\text{kg})(9.8\text{m/s}^2)$$

$$= 2822\text{ N}$$

$$F_{\text{net}} = F_{\text{app}} - F_f = ma$$

$$F_{\text{app}} = ma + F_f$$

$$= (1200)(5.85) + 2822$$

$$= \boxed{9800\text{ N}}$$

$$F_g = mg$$


on Earth's surface
 $\approx 9.8\text{m/s}^2$

$$F_f = \mu F_N$$

coefficient of friction

Normal Force supports \perp to surface

Ex: A boy pulls his 8.0 kg toboggan by a rope that angles 32° above the horizontal. If his 36.0 kg sister sits on the toboggan, how much force does he need to exert to accelerate them at 2.25 m/s^2 ? Assume $\mu=0$



$$F_{\text{net}} = F_x = ma$$


$$= (36.0 + 8.0)(2.25)$$

$$= 99 \text{ N}$$

$$\cos 32^\circ = \frac{F_x}{F_{\text{app}}}$$

$$F_{\text{app}} = \frac{F_x}{\cos 32^\circ} = \boxed{120 \text{ N}}$$

Ex: A 1.12 kg textbook is pushed horizontally against a wall with a coefficient of friction of 0.465. What is the least amount of force required to keep the book from slipping?



$$F_f = F_g$$

$$= mg$$

$$= (1.12 \text{ kg})(9.8 \text{ m/s}^2)$$

$$= 10.976 \text{ N}$$


$$F_f = \mu F_N \quad F_N = F_{\text{app}}$$

$$F_f = \mu F_{\text{app}}$$

$$F_{\text{app}} = \frac{F_f}{\mu} = \frac{10.976 \text{ N}}{0.465}$$

$$= \boxed{23.6 \text{ N}}$$

Ex: Two blocks ($m_1 = 2.0 \text{ kg}$ and $m_2 = 3.0 \text{ kg}$) are connected by a rope as shown. m_1 is pulled to the right with a force of 18 N. What is the tension in the rope connecting the two masses? Assume $\mu=0$



To find \vec{a} look at the entire system:

$$F_{\text{net}} = F_{\text{app}} = m_{\text{total}} a$$

$$a = \frac{F_{\text{app}}}{m_{\text{total}}} = \frac{18 \text{ N}}{(2.0 + 3.0)} = 3.6 \text{ m/s}^2$$

To find tension look at only 1 mass:

$$m_1 \quad \left| \quad m_2\right.$$

$$F_{\text{net}} = F_{\text{app}} - T = m_1 a \quad \left| \quad F_{\text{net}} = T = m_2 a\right.$$

$$T = F_{\text{app}} - m_1 a$$

$$= 18 - (2.0)(3.6)$$

$$= 10.8$$

$$= \boxed{11 \text{ N}}$$


$$T = (3.0)(3.6)$$

$$= 10.8$$

$$= \boxed{11 \text{ N}}$$

Ex: A 65 kg student stands on a bathroom scale in an elevator and notices that it reads 520 N.

Determine the magnitude and direction of the acceleration of the elevator.



$$F_g = mg$$

$$= (65 \text{ kg})(9.8 \text{ m/s}^2)$$

$$= 637 \text{ N}$$

Since $F_g > F_N$ it must be winning, so...

$$F_{\text{net}} = F_g - F_N = ma$$

$$a = \frac{F_g - F_N}{m} = \frac{637 - 520}{65}$$

$$= \boxed{1.8 \text{ m/s}^2 \text{ down}}$$